

When Prescribing

Epiduo[®] Forte (adapalene and benzoyl peroxide) Gel, 0.3%/2.5%



Only you can determine the right treatment for your patients. To ensure your patients receive Epiduo Forte Gel as prescribed, follow these steps:

- 1 Check "Dispense as Written"**
as there is no generic substitute for Epiduo Forte Gel
- 2 Include ICD-10 Codes**
and previous treatment failures in notes to pharmacist
- 3 Save as a Favorite**
in the EMR system to ensure the process is easier in the future
- 4 Remind Patients to Use myForte**
mobile app as part of their daily regimen
- 5 Download Patient Savings Card[†]**
and give to all commercially insured patients



Epiduo Forte Gel is now offered at **\$0 co-pay** for some patients with the Galderma CareConnect Card.*

The Galderma CareConnect Patient Savings Program[†] simply isn't like other programs. The prescriptions you write can be easily filled by pharmacies and your commercially-insured patients can pay less out-of-pocket. Easy enough?

For up to a 3-month supply of Epiduo Forte Gel:

- Commercially insured patients pay \$0
- Uninsured patients pay \$75
- No activation is required and the Galderma CareConnect Card can be used at any participating pharmacy

Learn more at GaldermaCC.com



*Galderma CareConnect is only available for commercially insured or uninsured patients. Patients who are enrolled in a government-run or government-sponsored healthcare plan with a pharmacy benefit are not eligible to use the Galderma CareConnect Patient Savings Card.

[†]Restrictions apply. Please see Galderma CareConnect Patient Savings Card for details.

Important Safety Information

Indication: Epiduo[®] Forte (adapalene and benzoyl peroxide) Gel, 0.3%/2.5% is indicated for the topical treatment of acne vulgaris. **Adverse Events:** In the pivotal study, the most commonly reported adverse reactions ($\geq 1\%$) in patients treated with Epiduo Forte Gel were skin irritation, eczema, atopic dermatitis and skin burning sensation.

Warnings/Precautions: Patients using Epiduo Forte Gel should avoid exposure to sunlight and sunlamps and wear sunscreen when sun exposure cannot be avoided. Erythema, scaling, dryness, stinging/burning, irritant and allergic contact dermatitis may occur with use of Epiduo Forte Gel and may necessitate discontinuation. When applying Epiduo Forte Gel, care should be taken to avoid the eyes, lips and mucous membranes.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.